

## Recombinant Human ECE1 Protein (His Tag)

**Catalog Number:** PKSH033691

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human ECE1 protein Gln90-Trp770, with an N-terminal His
<b>Calculated MW</b>	78.8 kDa
<b>Observed MW</b>	94-130 kDa
<b>Accession</b>	P42892
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Endothelin-Converting Enzyme-1 (ECE-1) is a single-pass type II transmembrane (TM) protein with a short cytoplasmic tail and a large ectodomain. ECE-1 is a zinc protease of the neprilysin (NEP) family, which also includes ECE-2, PEX, XCE, DINE, and Kell, and several NEP-like proteins. It is widely expressed and has several alternatively spliced forms that differ in their TM domain or cytoplasmic tail. All isoforms of ECE-1 are expressed in umbilical vein endothelial cells, polynuclear neutrophils, fibroblasts, atrium cardiomyocytes and ventricles. Endothelin-converting enzyme-1 is involved in the proteolytic processing of Endothelin-1 (EDN1), Endothelin-2 (EDN2), and Endothelin-3 (EDN3) to biologically active peptides. Defects in ECE1 are a cause of Hirschsprung disease, cardiac defects and autonomic dysfunction (HSCRCDAD). It is a form of Hirschsprung disease with skip-lesions defects, craniofacial abnormalities and other dysmorphic features, and autonomic dysfunction.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017