

Recombinant Human CNTFR/CNTFR-alpha Protein (His Tag)

Catalog Number: PKSH031308

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	Baculovirus-Insect Cells-derived Human CNTFR/CNTFR-alpha protein Met 1-Pro 346, with an C-terminal His
Calculated MW	36.0 kDa
Observed MW	45-48 kDa
Accession	NP_001833.1
Bio-activity	Immobilized human CNTFR at 10 µg/ml (100 µl/well) can bind biotinylated human CNTF with a linear ranger of 1. 28-160 ng/ml.

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 50mM Tris, 100mM NaCl, pH 8.0 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Ciliary neurotrophic factor(CNTF) is a member of the cytokine family. It is a polypeptide hormone that have functions in promoting neurotransmitter synthesis and neurite outgrowth in certain neuronal populations. It's actions appear to be restricted to the nervous system. Ciliary neurotrophic factor(CNTF) has biological effects through the activation of a multi-subunit receptor complex, consisting of an extracellular CNTF binding subunit(CNTF α ;) and two transmembrane signal transduction proteins: glycoprotein gp130 and LIF receptor. CNTF is considered as a potent survival factor of neurons and oligodendrocytes and may be relevant in reducing tissue destruction during inflammatory attacks. CNTF is also a survival factor for neurons of the peripheral sensory sympathetic, and ciliary ganglia. It has been reported that CNTF could be an agent that has therapeutic potential and possibly induces differentiation of large multipolar ganglionic phenotype in a subset of progenitors.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017