

Recombinant Cpn10 Monoclonal Antibody

catalog number: **AN301498L**

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human;Rat;Mouse
Immunogen	Recombinant human Cpn10 fragment
Host	Rabbit
Isotype	IgG, κ
Clone	A193
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution


WB	1:500-1:1000
IHC	1:200-1:1000
IF	1:50

Data




Western Blot with Cpn10 Monoclonal Antibody at dilution of 1:1000. Lane 1: MCF-7, Lane 2: 293, Lane 3: 4T1, Lane 4: PC-12


Observed-MW:11 kDa
Calculated-MW:11 kDa



Immunohistochemistry of paraffin-embedded Rat colon using Cpn10 Monoclonal Antibody at dilution of 1:100.



Immunohistochemistry of paraffin-embedded Human colon using Cpn10 Monoclonal Antibody at dilution of 1:100.



Immunofluorescent analysis of (100% Ice-cold methanol) fixed HeLa cells using anti-Cpn10 Monoclonal Antibody at dilution of 1:50.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.0

Co-chaperonin implicated in mitochondrial protein import and macromolecular assembly. Together with Hsp60, facilitates the correct folding of imported proteins. May also prevent misfolding and promote the refolding and proper assembly of unfolded polypeptides generated under stress conditions in the mitochondrial matrix. The functional units of these chaperonins consist of heptameric rings of the large subunit Hsp60, which function as a back-to-back double ring. In a cyclic reaction, Hsp60 ring complexes bind one unfolded substrate protein per ring, followed by the binding of ATP and association with 2 heptameric rings of the co-chaperonin Hsp10. This leads to sequestration of the substrate protein in the inner cavity of Hsp60 where, for a certain period of time, it can fold undisturbed by other cell components. Synchronous hydrolysis of ATP in all Hsp60 subunits results in the dissociation of the chaperonin rings and the release of ADP and the folded substrate protein.