Recombinant Human JAM-B/CD322 Protein (Fc Tag)

Catalog Number: PKSH032663

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Species	Human		
Source	HEK293 Cells-derived Human JAM-B;CD322 protein Phe29-Asn236, with an C-		
	terminal Fc		
Calculated MW	50.4 kDa		
Observed MW	60-75 kDa		
Accession	P57087		
Bio-activity	Not validated for activity		
Properties			
Purity	> 95 % as determined by reducing SDS-PAGE.		
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.		
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80		
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
	reconstituted samples are stable at $< -20^{\circ}C$ for 3 months.		
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.		
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants		
	before lyophilization.		
	Please refer to the specific buffer information in the printed manual.		
Reconstitution	Please refer to the printed manual for detailed information.		

Data

kDa	MK	R
120 90 60 40		ence
30)]	science
20 16 14	11	

> 95 % as determined by reducing SDS-PAGE.

Background

Junctional Adhesion Molecule B (JAM-B) is a single-pass type I membrane protein that belongs to the juctional adhesion molecules family. JAM-B includes a signal sequence (aa 1-28); an extracellular region (aa 29-238) with one Ig-like C2-type domain and one Ig-like V-type domain; a transmembrane segment (aa 239-259); and a cytoplasmic domain (aa 260 - 298). JAMB is localized to the tight junctions between endothelial cells or epithelial cells. JAM-B is prominently expressed in the heart; placenta; lung; foreskin and lymph node. It is also present on the endothelia of other vessels. JAM-B acts as an adhesive ligand for interacting with a variety of immune cell types and may play a role in lymphocyte homing to secondary lymphoid organs.

For Research Use Only

Tel:400-999-2100