Recombinant Human AMIGO2 Protein (Fc Tag)

Catalog Number: PKSH033770



| Description | | | | |
|----------------|------------------------------------------------------------------------------------------|--|--|--|
| Species | Human | | | |
| Mol_Mass | 67.5 kDa | | | |
| Accession | Q86SJ2 | | | |
| Bio-activity | Not validated for activity | | | |
| Properties | | | | |
| Purity | > 95 % as determined by reducing SDS-PAGE. | | | |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. | | | |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 | | | |
| | °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of | | | |
| | reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. | | | |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. | | | |
| Formulation | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. | | | |
| | Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants | | | |
| | before lyophilization. | | | |
| | Please refer to the specific buffer information in the printed manual. | | | |
| Reconstitution | Please refer to the printed manual for detailed information. | | | |

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| ata | | | |
|-----|-----------|-----------|---|
| | kDa | МК | R |
| | | | |
| | 120 90 | | |
| | 90 60 | | |
| | | | |
| | 40 30 | maneria : | |
| | | | |
| | 20 14 | | |
| | | | |

> 95 % as determined by reducing SDS-PAGE.

Background

Amphoterin-Induced Protein 2 (AMIGO2) is a single-pass type I membrane protein which belongs to the AMIGO family of immunoglobulin superfamily. Mature AMIGO2 contains an Ig-like C2-type (immunoglobulin-like) domain; 6 LRR (leucine-rich) repeats; a LRRCT domain; as well as a LRRNT domain. AMIGO2 is mainly expressed in in breast; ovary; cervix; and uterus; although lower in lung; colon; and rectum. AMIGO2 required for depolarization-dependent survival of cultured cerebellar granule neurons. AMIGO2 may mediate homophilic as well as heterophilic cell-cell interaction with AMIGO1 or AMIGO3. AMIGO2 may contribute to signal transduction through its intracellular domain; and may be required for tumorigenesis of a subset of gastric adenocarcinomas.

For Research Use Only