



A Reliable Research Partner in Life Science and Medicine

PerCP Anti-Mouse CD11c Antibody[N418]

Catalog Number: E-AB-F0991F

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse

Host Armenian Hamster Isotype Armenian Hamster IgG

Clone No. N418

Isotype Control [Product E-AB-F09852F]

Conjugation PerCP

Conjugation Information PerCP is designed to be excited by the blue laser (488 nm) and detected using an optical

filter centered near 675 nm (e.g., a 690/50 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.


Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Data



C57BL/6 murine splenocytes are stained with PerCP Anti-Mouse CD11c Antibody and FITC Anti-Mouse MHC II (I-A/I-E) Antibody (Left). Splenocytes stained with FITC Anti-Mouse MHC II (I-A/I-E) Antibody (Right) are used as control.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8 $^{\circ}\text{C}$ for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD11 antigen-like family member C;CD11c;Integrin alpha-X;Itgax;Leukocyte adhesion

Web: www.elabscience.cn

receptor p150+95

 Uniprot ID
 Q9QXH4

 Gene ID
 16411

For Research Use Only



Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Background

CD11c is a 150 kD glycoprotein also known as αX integrin, CR4, and p150. CD11c forms a $\alpha X\beta2$ heterodimer with $\beta2$ integrin (CD18). It is primarily expressed on dendritic cells, NK cells, a subset of intestinal intraepithelial lymphocytes (IEL), and some activated T cells. The $\alpha X\beta2$ integrin plays an important role in cell-cell contact by binding its ligands: iC3b, fibrinogen and CD54.