

Purified Anti-Mouse NKG2A/C/E Antibody[20D5]

catalog number: **AN004090P**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse
Immunogen	Recombinant Mouse NKG2A/C/E protein
Host	Rat
Isotype	Rat IgG2a, κ
Clone	20D5
Purification	>98%, Protein A/G purified
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications

Recommended Dilution

FCM	2 $\mu\text{g/mL}$ (1×10^5 - 5×10^5 cells)
------------	---

Data



C57/BL6 Mouse splenocytes were stained with 0.2 μg Purified Anti-Mouse NKG2A/C/E Antibody[20D5] (Right) and 0.2 μg Rat IgG2a, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Rat IgG Secondary Antibody, then anti-Mouse NK1.1 PE-conjugated Monoclonal Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

The NKG2 molecules are a family of lectin-like receptors that form heterodimers with CD94. NKG2/CD94 heterodimer are primarily expressed on NK cells, and a subset of CD8+ T cells. Studies of CD94/NKG2 heterodimers on NK cells have demonstrated that the NKG2 components transduce signals after ligand binding. NKG2A transduces inhibitory signals, while NKG2C and NKG2E transduce activating signals.

For Research Use Only