

## PE/Cyanine7 Anti-Human CD195/CCR5 Antibody[HEK/1/85a]

Catalog Number: E-AB-F1392H

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


Reactivity	Human
Host	Mouse
Isotype	Rat IgG2a, $\kappa$
Clone No.	HEK/1/85a
Isotype Control	PE/Cyanine7 Rat IgG2a, $\kappa$ Isotype Control[2A3] [Product E-AB-F09832H]
Conjugation	PE/Cyanine 7
Conjugation Information	PE/Cyanine7 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 775 nm (e.g., a 780/60 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

### Applications

### Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. <b>The amount of the reagent is suggested to be used 5 <math>\mu</math>L of antibody per test (million cells in 100 <math>\mu</math>L staining volume or per 100 <math>\mu</math>L of whole blood).</b> Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
-----	--

### Data



Staining of normal human peripheral blood cells with FITC Anti-Human CD3 Antibody and PE/Cyanine7 Anti-Human CD195 Antibody[HEK/1/85a/7a] (left) or PE/Cyanine7 Rat IgG2b,  $\kappa$  Isotype Control (right). Cells in the lymphocytes gate were used for analysis.

### Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

### Antigen Information

Alternate Names	CCR5;C-C chemokine receptor type 5;HIV-1 fusion co-receptor
-----------------	---

### For Research Use Only

**Uniprot ID**

P51681

**Gene ID**

1234

**Background**

CD195, also known as CCR5, is a 45 kD G protein-coupled seven transmembrane C C-chemokine receptor. It binds to MIP-1 $\alpha$ , MIP-1 $\beta$ , and RANTES and is expressed on a subset of T cells and monocytes. CD195 mediates an intracellular signal thought to induce cell differentiation and proliferation. CCR5 has also been shown to act as a co-receptor for R5 HIV-1 cell entry; modification of CCR5 by sulfation contributes to the efficiency of HIV-1 entry. Recent studies have shown CCR5 to play a role in a variety of other human diseases, ranging from infectious and inflammatory diseases to cancer.

**For Research Use Only**