



A Reliable Research Partner in Life Science and Medicine

Human IFN-β Antibody Pair Set

Catalog No. E-KAB-0035 Applications ELISA

Synonyms IFNB1, IFB, IFF, IFNB

Kit components & Storage

Title	Specifications	Storage
Human IFN-β Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Human IFN-β Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0035)		
		Human IFN-β Capture Antibody	Human IFN-β Detection Antibody (Biotin)	
Immunogen	Immunogen	Recombinant Human IFN-β protein	Recombinant Human IFN-β protein	
Information Swissprot		P01574		
Product details	Reactivity	Human	Human	
	Host	Rabbit	Rabbit	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%	
		glycerol, pH 7.4	protective protein, 50% glycerol, pH	
			7.4	
	Purify	Antigen Affinity	Antigen Affinity	
	Specificity	Detects Human IFN-β in ELISAs.		

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com



A Reliable Research Partner in Life Science and Medicine

Applications

Human IFN-β Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4μg/mL	Human IFN-β Capture Antibody	
Capture			Aist
ELISA Detection	1:1000-1:10000	Human IFN-β Detection Antibody (Biotin)	Nigor Depth O 0.11 1000 1000 10000 10000 Human IFN-β concentration(pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

This gene encodes a cytokine that belongs to the interferon family of signaling proteins, which are released as part of the innate immune response to pathogens. The protein encoded by this gene belongs to the type I class of interferons, which are important for defense against viral infections. In addition, type I interferons are involved in cell differentiation and anti-tumor defenses. Following secretion in response to a pathogen, type I interferons bind a homologous receptor complex and induce transcription of genes such as those encoding inflammatory cytokines and chemokines. Overactivation of type I interferon secretion is linked to autoimmune diseases. Mice deficient for this gene display several phenotypes including defects in B cell maturation and increased susceptibility to viral infection.

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com