

## APC Anti-Human CD5 Antibody[5D7]

Catalog Number: E-AB-F1371E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description


Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1, κ
Clone No.	5D7
Isotype Control	APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792E]
Conjugation	APC
Conjugation Information	APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

### Applications

### Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. <b>The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).</b> Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
-----	---

### Data



Human peripheral blood lymphocytes are stained with FITC Anti-human CD19 Antibody and APC Anti-Human CD5 Antibody[5D7] (Left). Lymphocytes are stained with FITC Anti-human CD19 Antibody and APC Mouse IgG1, κ Isotype Control (Right).

### Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Do not freeze.
Shipping	Ice bag

### Antigen Information

Alternate Names	LEU1;Leu-1;Ly-1;LymphocyteantigenT1/Leu-1;T-cellsurfaceglycoproteinCD5;T1;Tp67
Uniprot ID	P06127
Gene ID	921

### For Research Use Only

## Background

CD5 is a 67 kD single chain type I glycoprotein also known as Leu-1, Ly-1 and T1. It is a member of the scavenger receptor superfamily found on T cells, thymocytes, B cell subsets, chronic B lymphocytic leukemia (B-Cells), and peripheral blood dendritic cells. CD5 modulates T and B cell receptor signaling, thymocyte maturation, and T-B cell interactions upon binding to ligands such as CD72.