

## Biotin Anti-Mouse CD11c Antibody[N418]

**Catalog Number:** E-AB-F0991B

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

<b>Reactivity</b>	Mouse
<b>Host</b>	Armenian Hamster
<b>Isotype</b>	Armenian Hamster IgG
<b>Clone No.</b>	N418
<b>Isotype Control</b>	Biotin Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09853B]
<b>Conjugation</b>	Biotin
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

### Applications

### Recommended usage

<b>FCM</b>	Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \mu\text{g}$ per $10^6$ cells in 100 $\mu\text{L}$ volume or 100 $\mu\text{L}$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
------------	---

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	CD11 antigen-like family member C;CD11c;Integrin alpha-X;Itgax;Leukocyte adhesion receptor p150+95
<b>Uniprot ID</b>	Q9QXH4
<b>Gene ID</b>	16411
<b>Background</b>	CD11c is a 150 kD glycoprotein also known as $\alpha\text{X}$ integrin, CR4, and p150. CD11c forms a $\alpha\text{X}\beta\text{2}$ heterodimer with $\beta\text{2}$ integrin (CD18). It is primarily expressed on dendritic cells, NK cells, a subset of intestinal intraepithelial lymphocytes (IEL), and some activated T cells. The $\alpha\text{X}\beta\text{2}$ integrin plays an important role in cell-cell contact by binding its ligands: iC3b, fibrinogen and CD54.

### For Research Use Only