

## APC Anti-Human/Mouse Integrin $\beta$ 7 Antibody[FIB21]

Catalog Number: AN00423E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Human;Mouse
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG2b, $\kappa$
<b>Clone No.</b>	FIB21
<b>Isotype Control</b>	APC Rat IgG2b, $\kappa$ Isotype Control[LTF-2] [Product E-AB-F09842E]
<b>Conjugation</b>	APC
<b>Conjugation Information</b>	APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

### Applications

### Recommended usage

**FCM** Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

### Data



Staining of normal human peripheral blood cells with FITC Anti-Human CD3 Antibody and APC Anti-Human/Mouse integrin  $\beta$ 7 Antibody[FIB21] (left) or APC Rat IgG2a Isotype Control (right). Cells in the lymphocytes gate were used for analysis.

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	$\beta$ 7 Integrin;integrin $\beta$ p;ITGB7
<b>Uniprot ID</b>	P26010;P26011
<b>Gene ID</b>	3695;16421

### For Research Use Only

## Background

Integrin  $\beta 7$  is a 130 kD glycoprotein also known as integrin  $\beta p$ . It is a member of the Ig superfamily. In association with integrin  $\alpha 4$  or  $\alpha E$  chain,  $\beta 7$  forms  $\alpha 4/\beta 7$  or  $\alpha E/\beta 7$  heterodimer.  $\alpha 4/\beta 7$  (CD49d/ $\beta 7$ , LPAM-1) is expressed on the majority of peripheral lymphocytes, on small subsets of thymocytes, and bone marrow progenitors. LPAM-1 binds to several ligands, VCAM-1, MAdCAM-1 and fibronectin, and is involved in lymphocyte adhesion and some hematopoietic progenitor cells migration.  $\alpha E/\beta 7$  (CD103/ $\beta 7$ ,  $\alpha IEL/\beta 7$ ) is expressed on intestinal intraepithelial lymphocytes (IEL), dendritic epidermal T cells, T regulatory cells, a subset of CD8+ T cells in lymph nodes and lamina propria. CD103/ $\beta 7$  complex is thought to play a role in lymphocyte retention via interaction with its ligand E-Cadherin.