

Recombinant Human CD10/Neprilysin Protein (His Tag)

Catalog Number: PKSH033583

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human CD10/Neprilysin protein Tyr52-Trp750, with an N-terminal His
Calculated MW	80.9 kDa
Observed MW	94-100 kDa
Accession	P08473
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Concentration	Subject to label value.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
Formulation	Supplied as a 0.2 µm filtered solution of PBS, pH7.4.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Neprilysin/CD10(NEP) is a zinc metallopeptidase expressed at the cell surface of a variety of cells. The functions is both as an endopeptidase with a thermolysin-like specificity and as a dipeptidyl-carboxypeptidase. NEP has been shown to be involved in the degradation of enkephalins in the mammalian brain and the inactivation of circulating atrial natriuretic peptide. NEP has also been identified as the common acute lymphocytic leukemia antigen (CALLA), and is expressed on the surface of lymphocytes in some disease states. These and other observations have resulted in considerable interest in NEP as a target for analgesics and antihypertensive drugs. NEP is also a major degrading enzyme of amyloid β peptide ($A\beta$) in the brain, indicating that down-regulation of NEP activity, which could be caused by aging, can contribute to the development of Alzheimer's disease by promoting $A\beta$ accumulation.

For Research Use Only