Elabscience®


HSP40-4 Polyclonal Antibody


catalog number: E-AB-15115

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|--|
| Reactivity | Human;Mouse;Rat |
| Immunogen | Recombinant protein of human DNAJA1 |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Affinity purification |
| Buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Recommended Dilution |
| WB | 1:500-1:2000 |
| IHC | 1:50-1:200 |


Data





Western Blot analysis of 231, A431, Raji, Jurkat and HepG2 Immunohistochemistry of paraffin-embedded Human thyroid cell using HSP40-4 Polyclonal Antibody at dilution of 1:500 cancer using HSP40-4 Polyclonal Antibody at dilution of

1:30



Immunohistochemistry of paraffin-embedded Human liver cancer using HSP40-4 Polyclonal Antibody at dilution of

1:30

| Preparation & Storage | |
|-----------------------|---|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the |
| | temperature recommended. |
| | |

Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:techsupport@elabscience.com Fax: 1-832-243-6017

Elabscience®

DnaJ-like proteins interact with HSP 70 molecular chaperones and function to facilitate protein folding and mitochondrial protein import. HSP 40-4, also known as HDJ2, is the human DnaJ homolog that functions as a co-chaperone with a cysteine-rich zinc finger domain. The cellular redox enzyme thioredoxin interacts with HSP 40-4, and oxidation and reduction reversibly regulate HSP 40-4 function in response to the changing redox states of the cell. The zinc finger domain of HSP 40-4 may act as a redox sensor of chaperone-mediated protein-folding machinery, since HSP 40-4 inactivation leads to the oxidation of cysteine thiols and a simultaneous release of coordinated zinc. Loss of the HSP 40-4 protein may be linked to severe defects in spermatogenesis that involve aberrant androgen signaling.

For Research Use Only