# **Elabscience**®


### Elab Fluor<sup>®</sup> 700 Anti-Mouse CD18 Antibody[M18/2]

Catalog Number: E-AB-F1113M1

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Mouse
Host	Rat
lsotype	Rat lgG2a, κ
Clone No.	M18/2
Isotype Control	Elab Fluor <sup>®</sup> 700 Rat IgG2a, к Isotype Control[2A3] [Product E-AB-F09832M1]
Conjugation	Elab Fluor <sup>®</sup> 700
Conjugation Information	Elab Fluor <sup>®</sup> 700 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 719 nm (e.g., a 725/40 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 $\mu$ L of antibody per test (million cells in 100 $\mu$ L staining volume or per 100 $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of C57BL/6 murine splenocytes with Elab Fluor<sup>®</sup> 700 Anti-Mouse CD18 Antibody[M18/2] (filled gray

histogram) or Elab Fluor<sup>®</sup> 700 Rat IgG2a,  $\kappa$  Isotype Control (empty black histogram). Total viable cells were used for analysis.

Preparation & Storag	ye
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	CD18;Cell surface adhesion glycoproteins LFA-1/CR3/p150+95 subunit beta;
	Complement receptor C3 subunit beta;Integrin beta-2;Itgb2
Uniprot ID	P11835

### For Research Use Only

# **Elabscience**®

Gene ID Background

#### 16414

CD18 is a 95 kD protein, also known as integrin  $\beta$ 2 subunit. It is expressed on all leukocytes. CD18, in association with integrin  $\alpha$  chain CD11a, CD11b, and CD11c forms LFA-1, Mac-1, and  $\alpha$ X $\beta$ 2, respectively, and plays an important role in leukocytes adhesion. The CD18 integrin complexes bind ICAM-1 (CD54), ICAM-2 (CD102), ICAM-3 (CD50), iC3b, and fibrinogen.