

Recombinant Mouse CD226/DNAM-1 Protein (His Tag)

Catalog Number: PKSM041233

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse CD226/DNAM-1 protein Glu19-Pro254, with an C-terminal His
Calculated MW	27.6 kDa
Observed MW	35-50 kDa
Accession	Q8K4F0
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Reconstitution	Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information.

Data



Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V3.6

Mouse DNAX accessory molecule-1(DNAM-1) is a type I transmembrane glycoprotein that belongs to the immunoglobulin superfamily. As an activating receptor, it interacts with the ligands CD155 and CD112, and activates natural killer (NK) cells via its immunoreceptor tyrosine-based activatory motif (ITAM). Mature mouse DNAM-1 has extracellular domain (ECD) that contains two Ig-like C2-set domains, and possesses a cytoplasmic region that contains motifs for binding PDZ domains. DNAM-1 is expressed on several lymphoid and myeloid cell types and interacts with CD155/PVR and Nectin-2/CD112. Ligation of DNAM-1 promotes the activation of NK cells, CD8+ T cells, and mast cells, induces dendritic cell maturation, initiates megakaryocyte and activated platelet adhesion to vascular endothelial cells, and stimulates monocyte extravasation. Conversely, it inhibits the formation of osteoclasts. Platelet-endothelium interactions that are mediated by DNAM-1 enable the metastasis of tumor cells to the lung.