

# Recombinant Human PDK-1 Protein (His Tag)

Catalog Number:PKSH030854



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	PDK1
<b>Species</b>	Human
<b>Expression Host</b>	Baculovirus-Insect Cells
<b>Sequence</b>	Ser 29-Ala 436
<b>Accession</b>	NP_002601.1
<b>Calculated Molecular Weight</b>	48.6 kDa
<b>Observed molecular weight</b>	45 kDa
<b>Tag</b>	N-His

## Properties

<b>Purity</b>	> 90 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 8.5, 10% glycerol Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 90 % as determined by reducing SDS-PAGE.

## Background

Pyruvate dehydrogenase kinase, isozyme 1, also known as [Pyruvate dehydrogenase [lipoamide]] kinase isozyme 1, mitochondrial and PDK1, is a member of the PDK / BCKDK protein kinase family. PDK-1 is expressed predominantly in the heart. It contains one histidine kinase domain. Pyruvate dehydrogenase kinase (PDK) isoforms are molecular switches that downregulate the pyruvate dehydrogenase complex (PDC) by reversible phosphorylation in mitochondria. An inhibitory effect of lipoic acid on PDKs would result in less phosphorylation of E1 and hence increased PDC activity. At least two isoenzymic forms of pyruvate dehydrogenase kinase ( PDK-1 and PDK-2 ) may be involved in the regulation of enzymatic activity of mammalian pyruvate dehydrogenase complex by phosphorylation. PDK-3 appears to have the highest specific activity among the three isoenzymes. PDK-1 inhibits the mitochondrial pyruvate dehydrogenase complex

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human PDK-1 Protein (His Tag)

Catalog Number: PKSH030854



by phosphorylation of the E1 alpha subunit, thus contributing to the regulation of glucose metabolism.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017