Elabscience®

Recombinant Mouse IL-28B protein(His Tag)

Catalog Number: PKSM041474

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|---------------------|--|
| Species | Mouse |
| Source | E.coli-derived Mouse IL-28B protein Asp 20-Val 193, with an C-terminal His |
| Calculated MW | 20.6 kDa |
| Observed MW | 17-25 kDa |
| Accession | Q8CGK6 |
| Bio-activity | Measure by its ability to protect HepG2 cells infected with encephalomyocarditis |
| | (EMC) virus. The ED ₅₀ for this effect is <20 ng/mL. |
| Properties | |
| Purity | > 98 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 0.1 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 |
| | °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of |
| | reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile PBS, pH 7.4. |
| | Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants |
| | before lyophilization. |
| | Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |
| Data | |
| k | Da |
| | |
| | 75- 63- |
| | 48- |
| | 35- |
| | 25- |
| | |
| | 17- |

> 98 % as determined by reducing SDS-PAGE.

11-

Background

Interleukin-28B; also known as Cytokine Zcyto22; Interferon lambda-3; Interferon lambda-4; IFNL3; IFNL4; ZCYTO22 and IL28B; is a secreted cytokine which belongs to the IL-28/IL-29 family. IL-28 has also been shown to play a role in the adaptive immune response. IL28B has immunomodulatory activity and up-regulates MHC class I antigen expression. IL28B displays potent antiviral activity and antitumor activity. In addition; IL28B is a ligand for the heterodimeric class II cytokine receptor composed of IL10RB and IL28RA. The ligand/receptor complex seems to signal through the Jak-STAT pathway.