Recombinant Human Interleukin-2/IL-2 Protein (His Tag)

Catalog Number: PKSH033425

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Species	Human	
Source	E.coli-derived Human Interleukin-2/IL-2 protein Ala21-Thr153, with an C-terminal His	
Calculated MW	16.4 kDa	
Observed MW	13 kDa	
Accession	P60568	
Bio-activity	Measure by its ability to induce proliferation in CTLL-2 cells. The ED_{50} for this effect	
	is <3 ng/mL. The specific activity of recombinant human IL-2 is >5 x 10^6 IU/mg.	
Properties		
Purity	> 98 % as determined by reducing SDS-PAGE.	
Endotoxin	< 0.01 EU per µg of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80	
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20 °C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from sterile PBS,pH 8.0.	
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants	
	before lyophilization.	
	Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	
Data		
	·····	

kDa	MK	R
120 90	-	
60	-	
40		
30	-	
20	-	a formation
	and the second	
14		-

> 98 % as determined by reducing SDS-PAGE.

Background

Interleukin-2(IL-2) is an interleukin; a type of cytokine signaling molecule in the immune system; belongs to the IL-2 family. It is a powerful immunoregulatory lymphokine produced by T-cells in response to antigenic or mitogenic stimulation. IL-2/IL-2R signaling is required for T-cell proliferation and other fundamental functions that are essential for the immune response. IL-2 stimulates growth and differentiation of B-cells; NK cells; lymphokine-activated killer cells; monocytes; macrophages and oligodendrocytes.

Web:www.elabscience.com