

Recombinant Rat SerpinF1/PEDF Protein (His Tag)

Catalog Number: PKSR030156

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Rat
Source	HEK293 Cells-derived Rat SerpinF1/PEDF protein Met1-Thr418, with an C-terminal His
Calculated MW	45.9 kDa
Accession	NP_808788.1
Bio-activity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Pigment epithelium-derived factor, also known as PEDF, Serpin F1, and SERPINF1, is a multiple functional protein which has both anti-angiogenic activity and neurotrophic activity at the same time. PEDF is a secreted glycoprotein that belongs to the noninhibitory serpin. It has an alpha/beta core serine-protease inhibitor domain, three major beta-sheets, and ten alpha-helices. PEDF does not inhibit either serine or cysteine proteinases. PEDF exerts diverse physiological activities including anti-angiogenesis, anti-vasopermeability, anti-tumor, and neurotrophic activities. PEDF acts via multiple high affinity ligands and cell receptors. It has been described as a natural angiogenesis inhibitor with neurotrophic and immune-modulation properties. PEDF induces macrophages apoptosis and necrosis through the activation of peroxisome proliferator-activated receptor-gamma by which PEDF could modulate inflammatory reactions in septic shock. It balances angiogenesis in the eye and blocks tumor progression.

For Research Use Only