

Recombinant Mouse IL-21 Protein(Sumo Tag)

Catalog Number: PDEM100122

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	E.coli-derived Mouse IL-21 protein His18-Ser146, with an N-terminal Sumo
Mol_Mass	27.0 kDa
Accession	Q9ES17
Bio-activity	Not validated for activity

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Mouse IL-21 proteins, 2 µg/lane of Recombinant Mouse IL-21 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 27.0 KD

Background

IL21 belongs to the IL-15/IL-21 family. It is a cytokine with immunoregulatory activity. Cytokines are proteinaceous signaling compounds that are major mediators of the immune response. They control many different cellular functions including proliferation, differentiation, and cell survival/apoptosis but are also involved in several pathophysiological processes including viral infections and autoimmune diseases. Cytokines are synthesized under various stimuli by a variety of cells of both the innate (monocytes, macrophages, dendritic cells) and adaptive (T-and B-cells) immune systems. IL21 is expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. It may promote the transition between innate and adaptive immunity. IL-21 has been tried as a therapy for alleviating allergic responses. It can significantly decrease pro-inflammatory cytokines produced by T cells in addition to decreasing IgE levels in a mouse model for rhinitis (nasal passage inflammation).

For Research Use Only