

Recombinant Rat Cadherin-8/CDH8 Protein (His Tag)

Catalog Number: PKSR030268

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Rat
Source	HEK293 Cells-derived Rat Cadherin-8/CDH8 protein Met1-Met621, with an C-terminal His
Calculated MW	66.5 kDa
Observed MW	75-85 kDa
Accession	NP_445845.2
Bio-activity	Not validated for activity

Properties

Purity	> 85 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 85 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Cadherins are integral membrane proteins that mediate calcium-dependent cell-cell adhesion. Type I cadherin proteins are composed of a large N-terminal extracellular domain, a single membrane-spanning domain, and a small, highly conserved C-terminal cytoplasmic domain. The extracellular domain consists of five subdomains, each containing a cadherin motif, and appears to determine the specificity of the protein's homophilic cell adhesion activity. Type II (atypical) cadherins are defined based on their lack of a HAV cell adhesion recognition sequence specific to type I cadherins. Cadherin 8, also known as CDH 8, is a type II classical cadherin belonging to the cadherin superfamily. As mainly expressed in brain, CDH8 is found in certain nerve cell lines, such as retinoblasts, glioma cells and neuroblasts, and is putatively involved in synaptic adhesion, axon outgrowth and guidance. Human Cadherin 8 is a 799 amino acid single-pass type I transmembrane protein with a putative 29 aa signal sequence, and a 32 aa propeptide, a 560 aa mature extracellular domain, a 21 aa transmembrane domain and a 157 aa cytoplasmic domain. The human, mouse and rat proteins share approximately 98% homology.