# **Elabscience**®

### Human MMP-2 Antibody Pair Set

Catalog No.E-KAB-0451ApplicationsSynonymsMMP2;CLG4;CLG4A;MMP-II;MONA;TBE-1;Gelatinase A

ELISA

#### Kit components & Storage

Title	Specifications	Storage
Human MMP-2 Capture Antibody	1 vial, 100 µ g	Store at -20°C for one year. Avoid
		freeze/thaw cycles.
Human MMP-2 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year. Avoid
(Biotin)		freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0451)	
		Human MMP-2 Capture Antibody	Human MMP-2 Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Human MMP-2 protien	Recombinant Human MMP-2 protien
Information	Swissprot	P08253	
Product details	Reactivity	Human	Human
	Host	Rabbit	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%
		glycerol; pH 7.5	protective protein; 50% glycerol; pH
			7.5
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Human MMP-2 in ELISAs.	

For Research Use Only

## **Elabscience**®

#### Applications

Human MMP-2 Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Human MMP-2 Capture	
Capture		Antibody	10
			Ag 1
			Optical Density
ELISA	1:1000-1:10000	Human MMP-2 Detection	O Obii
Detection		Antibody (Biotin)	0.01
			0.01 1 10 100
			Human MMP-2 Concentration(ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

#### Background

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. This gene encodes an enzyme which degrades type IV collagen, the major structural component of basement membranes. The enzyme plays a role in endometrial menstrual breakdown, regulation of vascularization and the inflammatory response.

For Research Use Only