

Recombinant Human Noggin/NOG Protein (His Tag)

Catalog Number: PKSH031720

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human Noggin/NOG protein Met 1-Cys 232, with an C-terminal His
Calculated MW	24.6 kDa
Observed MW	30 kDa
Accession	NP_005441.1
Bio-activity	Measured by its ability to inhibit recombinant human BMP4-induced alkaline phosphatase production by MC3T3-E1 cells. The ED ₅₀ for this effect is typically 0.05-0.3 µg/mL in the presence of 50 ng/mL of recombinant human BMP4.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



Measured by its ability to inhibit recombinant human BMP4-induced alkaline phosphatase production by MC3T3-E1 cells. The ED₅₀ for this effect is typically 10-60 ng/mL in the presence of 25 ng/mL of recombinant human BMP4.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Noggin is a secreted protein involved at multiple stages of vertebrate embryonic development including neural induction and is known to exert its effects by inhibiting the bone morphogenetic protein (BMP)-signaling pathway. It binds several BMPs with very high (picomolar) affinities; with a marked preference for BMP2 and BMP4 over BMP7. By binding tightly to BMPs; Noggin prevents BMPs from binding their receptors. Noggin binds the bone morphogenetic proteins (BMP) such as BMP-4 and BMP-7; and inhibits BMP signaling by blocking the molecular interfaces of the binding epitopes for both type I and type II receptors. Interaction of BMP and its antagonist Noggin governs various developmental and cellular processes; including embryonic dorsal-ventral axis; induction of neural tissue; formation of joints in the skeletal system and neurogenesis in the adult brain. Noggin plays a key role in neural induction by inhibiting BMP4; along with other TGF- β ; signaling inhibitors such as chordin and follistatin. Mouse knockout experiments have demonstrated that noggin also plays a crucial role in bone development; joint formation; and neural tube fusion.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017