

AF/LE Purified Anti-Mouse Ly-6G/Ly-6C (Gr-1) Antibody[RB6-8C5]

catalog number: E-AB-F11200



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


| | |
|---------------------|--|
| Reactivity | Mouse |
| Immunogen | Recombinant Mouse Gr-1 protein |
| Host | Rat |
| Isotype | Rat IgG2b, κ |
| Clone | RB6-8C5 |
| Purification | >98%, Protein A/G purified |
| Conjugation | None (AF/LE) |
| buffer | Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method |

Applications

Recommended Dilution

| | |
|------------|---|
| FCM | 2 $\mu\text{g/mL}$ (1×10^5 - 5×10^5 cells) |
|------------|---|

Data



C57BL/6 Mouse bone marrow cells were stained with 0.2 μg AF/LE Purified Anti-Mouse Ly-6G/Ly-6C (Gr-1) Antibody[RB6-8C5] (Right) and 0.2 μg Rat IgG2b, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Rat IgG Secondary Antibody.

Preparation & Storage

| | |
|-----------------|--|
| Storage | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions. |
| Shipping | Ice bag |

Background

Gr-1 is a 21-25 kD protein also known as Ly-6G/Ly-6C. This myeloid differentiation antigen is a glycosylphosphatidylinositol (GPI)-linked protein expressed on granulocytes and macrophages. In bone marrow, the expression levels of Gr-1 directly correlate with granulocyte differentiation and maturation; Gr-1 is also transiently expressed on bone marrow cells in the monocyte lineage. Immature Myeloid Gr-1+ cells play a role in the development of antitumor immunity.

For Research Use Only