A Reliable Research Partner in Life Science and Medicine

# PE/Cyanine5 Anti-Mouse CD127/IL-7RA Antibody[A7R34]

Catalog Number: E-AB-F1023G

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Reactivity Mouse Host Rat

**Isotype** Rat IgG2a, κ **Clone No.** A7R34

Isotype Control PE/Cyanine5 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832G]

Conjugation PE/Cyanine 5

Conjugation Information PE/Cyanine5 is designed to be excited by the Blue (488 nm), Green (532 nm) and

yellow-green (561 nm) lasers and detected using an optical filter centered near 670 nm

(e.g., a 690/50 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


### Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

# Data



C57BL/6 murine splenocytes are stained with PE/Cyanine5

Anti-Mouse CD127/IL-7RA Antibody and Elab Fluor<sup>®</sup> 488 Anti-Mouse CD3 Antibody (Left). Splenocytes stained with

Elab Fluor® 488 Anti-Mouse CD3 Antibody (Right) are used as control.

## **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

#### **Antigen Information**

Alternate Names CD127;IL-7 receptor subunit alpha;IL-7R subunit alpha;IL-7R-alpha;IL-7RA;II7r;

Interleukin-7 receptor subunit alpha

# For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web:www.elabscience.com
 Email:techsupport@elabscience.com

# Elabscience®

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

**Uniprot ID** P16872 Gene ID 16197 **Background** 

CD127 is a 60-90 kD type I transmembrane glycoprotein also known as IL-7 receptor  $\alpha$ chain or IL-7Rα. It forms a heterodimer with the common γ chain (γc or CD132) which is shared with the receptors for IL-2, IL-4, IL-9, IL-13, IL-15, and IL-21. CD127 is expressed on immature B cells through early pre-B stage, thymocytes (except CD4/ CD8 double positive thymocytes), peripheral T cells, and bone marrow stromal cells. CD127 has been reported to be an useful marker for identifying memory and effector T cells. The ligation of IL-7 with its receptor is important for stimulation of mature and immature T cells as well as immature B cells proliferation and development.

Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web:www.elabscience.com Email:techsupport@elabscience.com