



A Reliable Research Partner in Life Science and Medicine

## Recombinant Human COL2A1 Protein (GST, His Tag)

Catalog Number: PDEH100914

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source E.coli-derived Human COL2A1 protein Met1-Leu268, with an N-terminal GST & C-

erminal His

 Calculated MW
 54.4 kDa

 Observed MW
 54 kDa

 Accession
 P02458-3

Bio-activity Not validated for activity

**Properties** 

**Purity** > 95% as determined by reducing SDS-PAGE.

Endotoxin < 10 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -

80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping

This product is provided as lyophilized powder which is shipped with ice packs.

Formulation

Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

**Reconstitution** It is recommended that sterile water be added to the vial to prepare a stock solution

of 0.5 mg/mL. Concentration is measured by UV-Vis.

## **Background**

COL2A1 is the alpha-1 chain of type II collagen which is a fibrillar collagen found in cartilage and the vitreous humor of the eye. Mutations in this protein are associated with achondrogenesis, chondrodysplasia, early onset familial osteoarthritis, SED congenita, Langer-Saldino achondrogenesis, Kniest dysplasia, Stickler syndrome type I, and spondyloepimetaphyseal dysplasia Strudwick type. In addition, defects in processing chondrocalcin, a calcium binding protein that is the C-propeptide of this collagen molecule, are also associated with chondrodysplasia. There are two transcripts identified for this gene. Type II collagen is specific for cartilaginous tissues. Thus COL2A1 is essential for the normal embryonic development of the skeleton, for linear growth and for the ability of cartilage to resist compressive forces. The regulation of COL2A1, likely results from a balance of both positive and negative proteins. The inhibition of COL2A1 transcription following treatment of chick sternal chondrocytes with growth factors was accompanied by increased EF1 expression. Overexpression of EF1 in differentiated chondrocytes resulted in decreased expression of a reporter construct containing a collagen II promoter/enhancer insert.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com