Recombinant SARS-CoV-2 Spike RBD-SD1(V367F)(His Tag)

Catalog Number: PKSV030386

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Species	SARS-CoV-2
Source	HEK293 Cells-derived SARS-CoV-2 SARS-CoV-2 Spike RBD-SD1(V367F) protein Arg319-Ser591(Val367Phe), with an C-terminal His
Calculated MW	141 kDa
Observed MW	30-40 kDa
Accession	QHD43416.1
Bio-activity	PKSR030477Immobilized Human ACE-2-Fc (PKSR030492) at 5 μg/ml (100 μl/well) can bind 2019-nCoV S Protein RBD-SD1-His (PKSR030477). The ED_{50} of PKSR030477 is 1.7 ug/ml. PKSV030386Immobilized Human ACE-2-Fc (PKSR030492) at 5 μg/ml (100 μl/well) can bind 2019-nCoV S Protein RBD-SD1(V367F, C-6His)(PKSV030386). The ED_{50} of PKSV030386 is 0.8 ug/ml., PKSR030477Immobilized Anti-2019-nCoV S1 mAb (5D9) at 5 μg/ml (100 μl/well) can bind 2019-nCoV S Protein RBD-SD1-His (PKSR030477). The ED_{50} of PKSR030477 is 1. 28 ug/ml. PKSV030386Immobilized Anti-2019-nCoV S1 mAb (5D9) at 5 μg/ml (100 μl/well) can bind 2019-nCoV S Protein RBD-SD1(V367F, C-6His)(PKSV030386). The ED_{50} of PKSV030386 is 2. 55 ug/ml.

Properties

Purity	> 95% as determined by reducing SDS-PAGE.
Concentration	Subject to label value.
Endotoxin	Please contact us for more information.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/ gel packs. Upon receipt, store it immediately at < - 20°C.
Formulation	Supplied as a 0.2 μm filtered solution of PBS, pH7.4.

Data



Background

The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cel l. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection proces s. It's been reported that SARS-CoV-2 (COVID-19 coronavirus, 2019-nCoV) can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity. The main functions for the Spike protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate.

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: w w w .elabscience.com
Email: techsupport@elabscience.com