

IBSP Polyclonal Antibody

catalog number: E-AB-19332

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|--------------|--|
| Reactivity | Human;Mouse |
| Immunogen | Synthetic peptide of human IBSP |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Antigen affinity purification |
| Conjugation | Unconjugated |
| buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |

Applications

| Applications | Recommended Dilution |
|--------------|----------------------|
| WB | 1:500-1:2000 |
| IHC | 1:25-1:100 |


Data



Western blot analysis of Jurkat cell lysate using IBSP Polyclonal Antibody at dilution of 1:700

Observed-MV:Refer to figures

Calculated-MV:35 kDa



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using IBSP Polyclonal Antibody at dilution of 1:30(×200)

Preparation & Storage

| | |
|----------|--|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended. |

Background

The protein encoded by this gene is a major structural protein of the bone matrix. It constitutes approximately 12% of the noncollagenous proteins in human bone and is synthesized by skeletal-associated cell types, including hypertrophic chondrocytes, osteoblasts, osteocytes, and osteoclasts. The only extraskeletal site of its synthesis is the trophoblast. This protein binds to calcium and hydroxyapatite via its acidic amino acid clusters, and mediates cell attachment through an RGD sequence that recognizes the vitronectin receptor.

For Research Use Only