

## Recombinant Human Interleukin-33/IL-33 Protein

Catalog Number: PKSH033616

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	E.coli-derived Human Interleukin-33/IL-33 protein Ser112-Thr270
<b>Calculated MW</b>	18.1 kDa
<b>Observed MW</b>	18 kDa
<b>Accession</b>	O95760
<b>Bio-activity</b>	Immobilized Mouse ST2-His(PKSM041050) at 5µg/ml (100 µl/well) can bind Human IL-33(PKSH033616).The ED <sub>50</sub> of Human IL-33(PKSH033616) is 0.63 µg/ml.


### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 0.01 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.



Immobilized Mouse ST2-His(PKSM041050) at 5µg/ml (100 µl/well) can bind Human IL-33(PKSH033616).The ED<sub>50</sub> of Human IL-33(PKSH033616) is 0.63 µg/ml.

### Background

#### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Interleukin-33 (IL-33) was initially discovered as a nuclear factor NF-HEV abundantly expressed in high endothelial venules. It is a 30-32 kD pro-inflammatory protein with intracellular and extracellular activities and a chromatin-associated cytokine of the IL-1 family with high sequence and structural similarity to IL-1 and IL-18. IL-33 is highly and selectively expressed by high endothelial venule endothelial cells (HEVECs) in human tonsils; Peyers's patches; and lymph nodes. It contains a bipartite nuclear localization signal at the C-terminus; and is targeted to the nucleus when ectopically expressed in human umbilical vein endothelial cells (HUVECs) and HeLa cells. The C-terminal fragment; corresponding to mature IL-33; binds and triggers signaling. IL-33 mediates its biological effects via Toll-interleukin 1 (IL-1) receptor (TIR) domain-containing receptor ST2; activates NF-kappaB and MAP kinases; and drives production of T(H)2-associated cytokines from in vitro polarized T(H)2 cells. In vivo; IL-33 induces the expression of IL-4; IL-5; and IL-13 and leads to severe pathological changes in mucosal organs. Human IL-33 is 270 amino acids in length.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017