

A Reliable Research Partner in Life Science and Medicine

PE/Cyanine5 Goat Anti-Mouse IgG (H+L) Antibody[Poly1440]

Catalog Number: AN00338G

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse **Host** Goat

Isotype Goat Polyclonal IgG

Clone No. Poly1440
Conjugation PE/Cyanine 5

Conjugation Information PE/Cyanine5 is designed to be excited by the Blue (488 nm), Green (532 nm) and

yellow-green (561 nm) lasers and detected using an optical filter centered near 670 nm

(e.g., a 690/50 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of normal human peripheral blood cells with purified mouse anti-human CD3, followed by PE/Cyanine5 Goat Anti-Mouse IgG (H+L) Antibody[Poly1440] (filled gray histogram). Cells in the lymphocytes gate were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Web: www.elabscience.cn

Shipping Ice bag

Antigen Information

Alternate Names Goat Anti-Mouse IgG

 Uniprot ID
 Q61559

 Gene ID
 16059

For Research Use Only



Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Background

This polyclonal goat anti-mouse IgG antibody reacts with the heavy chains of mouse IgG and with the light (kappa and lambda) chains common to most mouse immunoglobulins. No cross-reactivity has been detected against non-immunoglobulin serum proteins. This antibody has been solid-phase absorbed to ensure minimal cross-reaction with rat, human, bovine, horse, and rabbit immunoglobulins, but it may have minimal cross-reactivity with other subclasses of mouse immunoglobulins.