

Recombinant Human CHGA protein (His tag)

Catalog Number:PDMH100095



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Chromogranin-A;CgA;Pituitary secretory protein I;SP-I;CHGA
Species	Human
Expression Host	HEK293 Cells
Sequence	Met1-Gly457
Accession	P10645
Calculated Molecular Weight	50.2 kDa
Observed molecular weight	75 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Chromogranin A (CgA), also known as pituitary secretory protein I (SP-I), is a member of the granin family of regulated secretory proteins. CgA shares several protein characteristics common to the granin family: acidic isoelectric point, the capacity to bind calcium ions, the ability to form aggregates and multiple dibasic cleavage sites. Mature human CgA is 439 amino acids (aa) and contains 10 dibasic, proteolytic cleavage sites, capable of yielding several smaller peptides, each displaying a unique function. Mature human CgA shares 63% aa sequence identity with mouse and rat CgA. CgA is expressed exclusively in the secretory dense core granules of most normal and neoplastic neuroendocrine cells. Increased levels of CgA have been detected in patients with neuroendocrine tumors as well as non-neuroendocrine tumors, hence CgA is an important serological marker for tumor diagnosis and monitoring tumor progression/regression. It has been

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human CHGA protein (His tag)

Catalog Number:PDMH100095



demonstrated in mouse model that full-length CgA containing its C-terminal region can impair angiogenesis and tumor growth. In addition, CgA can bind to Secretogranin III to regulate the biogenesis of secretory granules.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017