

Human IL-12 Antibody Pair Set

Catalog No.	E-KAB-0477	Applications	ELISA
Synonyms	IL12;IL-12/p70;IL12 p70		

Kit components & Storage

Title	Specifications	Storage
Human IL-12 Capture Antibody	1 vial, 100 µg	Store at -20℃ for one year. Avoid freeze/thaw cycles.
Human IL-12 Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20℃ for one year. Avoid freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Product Information

Items		Characteristic (E-KAB-0477)	
		Human IL-12 Capture Antibody	Human IL-12 Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Human IL-12 protien	Recombinant Human IL-12 protien
	Swissprot	P29459	
Product details	Reactivity	Human	Human
	Host	Rat	Mouse
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50% glycerol; pH 7.5	PBS with 0.04% Proclin 300; 1% protective protein; 50% glycerol; pH 7.5
	Purify	Protein A or G	Protein A or G
	Specificity	Detects Human IL-12 in ELISAs.	

For Research Use Only

Applications

Human IL-12 Sandwich ELISA Assay

	Recommended Concentration/Dilution	Reagent	Images
ELISA Capture	0.5-4 µg/mL	Human IL-12 Capture Antibody	
ELISA Detection	1:1000-1:10000	Human IL-12 Detection Antibody (Biotin)	

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

This gene encodes a subunit of a cytokine that acts on T and natural killer cells , and has a broad array of biological activities. The cytokine is a disulfide-linked heterodimer composed of the 35-kD subunit encoded by this gene , and a 40-kD subunit that is a member of the cytokine receptor family. This cytokine is required for the T-cell-independent induction of interferon (IFN) -gamma , and is important for the differentiation of both Th1 and Th2 cells. The responses of lymphocytes to this cytokine are mediated by the activator of transcription protein STAT4. Nitric oxide synthase 2A (NOS2A/NOS2) is found to be required for the signaling process of this cytokine in innate immunity.

For Research Use Only