

## Recombinant Human JAM-A/F11R Protein (His Tag)

**Catalog Number:** PKSH031771

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human JAM-A/F11R protein Met 1-Ala 242, with an C-terminal His
<b>Mol_Mass</b>	25 kDa
<b>Accession</b>	NP_058642.1
<b>Bio-activity</b>	Measured by the ability of the immobilized protein to support the adhesion of Jurkat human acute T cell leukemia cells. When $8 \times 10^4$ cells/well are added to JAM-A-Fc coated plates (2.5 µg/mL, 100 µL/well) in the presence of 20 ng/mL PMA, approximately 30-40% will adhere after 30 minutes at 37°C.

### Properties

<b>Purity</b>	> 97 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 97 % as determined by reducing SDS-PAGE.

### Background

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Junctional adhesion molecule-A (JAM-A); also known as F11 receptor (F11R) or Cluster of Differentiation 321 (CD321); is a transmembrane protein expressed at tight junctions of epithelial and endothelial cells; as well as on circulating leukocytes. JAM-A protein serves as a serotype-independent receptor for mammalian orthoreoviruses (reoviruses). It is also a ligand for the integrin LFA1; involves in leukocyte transmigration. As a cell adhesion molecule of the immunoglobulin superfamily; JAM-A protein involves in platelet adhesion; secretion and aggregation; and plays a crucial role in inflammatory thrombosis and atherosclerosis. In addition; it may be a potential therapeutic target for breast cancer.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V3.4