

A Reliable Research Partner in Life Science and Medicine

FITC Anti-Mouse CD122/IL-2RB Antibody[5H4]

Catalog Number: E-AB-F1029UC

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse **Host** Rat

lsotype Rat lgG2a, κ

Clone No. 5H4

Isotype Control FITC Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833C]

Conjugation FITC

Conjugation Information FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical

filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/10⁶ cells in 100 μ L volume].

Data



C57BL/6 murine splenocytes are stained with APC Anti-Mouse NK1.1 Antibody and FITC Anti-Mouse CD122 Antibody (Left). Splenocytes are stained with APC Anti-Mouse NK1.1 Antibody and FITC Rat IgG2a, κ Isotype Control (Right).

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names IL-2R subunit beta;CD122;High affinity IL-2 receptor subunit beta;IL-2 receptor subunit

beta;IL-2RB;II2rb;Interleukin-2 receptor subunit beta;p70-75

Uniprot ID P16297

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Gene ID Background 16185

CD122 is a 70-75 kD IL-2 receptor β chain also known as IL-2R β , which is also shared by the IL-15 receptor. It is constitutively expressed by NK cells and at lower levels by T cells, B cells, monocytes, and macrophages. The IL-2R β chain can combine with either the common γ subunit (γ c, CD132) alone or with the γ c subunit and the IL-2R β subunit (CD25) to generate intermediate or high affinity IL-2 receptor complexes, respectively. CD122 expression levels can be upregulated by activation. The 5H4 antibody does not block IL-2 binding to the IL-2 receptor. CD122 is expressed on murine, but not human, CD8+ Tregs involved in the maintenance of T cell homeostasis.