

Recombinant Mouse TIM4/TIMD4 Protein (His Tag)

Catalog Number:PKSM040707



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	B430010N18Rik;TIM-4;Tim4
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Thr 279
Accession	NP_848874.3
Calculated Molecular Weight	29.3 kDa
Observed molecular weight	60-70 kDa
Tag	C-His

Properties

Purity	> 94 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 94 % as determined by reducing SDS-PAGE.

Background

A type I transmembrane protein called TIM4 (T-cell immunoglobulin- and mucin-domain-containing molecule; also known as TIMD4), which belongs to the immunoglobulin superfamily and TIM family. TIM4 is involved in regulating T-cell proliferation and lymphotoxin signaling. It is a ligand for HAVCR1/TIMD1. Recent reports indicate that dendritic cell (DC)-derived T-cell immunoglobulin and mucin domain molecule (TIM)-4, which is expressed on dendritic cells and macrophages, plays an important role in the initiation of T(H)2 polarization. TIM4 bound apoptotic cells by recognizing phosphatidylserine via its immunoglobulin domain. The expression of TIM4 in fibroblasts enhanced their ability to engulf apoptotic cells. TIM4 is phosphatidylserine receptor for the engulfment of apoptotic cells, and may also be involved in intercellular signalling in which exosomes are involved. Modulation of TIM4 production in dendritic cells (DCs)

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Mouse TIM4/TIMD4 Protein (His Tag)

Catalog Number:PKSM040707



represents a novel therapeutic approach for the treatment of peanut allergy. The interaction of TIM1/TIM4 played a critical role in sustaining the polarization status of Th2 cells in allergic rhinitis (AR) patients. Cross-linking FcγRI by antigen/IgG complexes increased the production of TIM4 by dendritic cells via upregulating tumor necrosis factor-α in DCs. Specific immunotherapy (SIT) suppresses the skewed Th2 responses via disrupting the interaction of TIM1/TIM4 in antigen-specific Th2 cells.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com