

## Elab Fluor® 700 Anti-Human/Monkey CD196/CCR6 Antibody[G034E3]

Catalog Number: E-AB-F1158M1

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


|                                |  |
|--------------------------------|--|
| <b>Reactivity</b>              | Human;Rhesus   |
| <b>Host</b>                    | Mouse  |
| <b>Isotype</b>                 | Mouse IgG2b, κ   |
| <b>Clone No.</b>               | G034E3   |
| <b>Isotype Control</b>         | Elab Fluor® 700 Mouse IgG2b, κ Isotype Control[MPC-11] [Product E-AB-F09812M1]   |
| <b>Conjugation</b>             | Elab Fluor® 700  |
| <b>Conjugation Information</b> | Elab Fluor® 700 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 719 nm (e.g., a 725/40 nm bandpass filter). |
| <b>Storage Buffer</b>          | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.  |

### Applications

### Recommended usage

|            |   |
|------------|---|
| <b>FCM</b> | Each lot of this antibody is quality control tested by flow cytometric analysis. <b>The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).</b> Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. |
|------------|---|

### Data



Staining of normal human peripheral blood cells with FITC

Anti-Human CD19 Antibody[CB19] and Elab Fluor® 700  
Anti-Human/Monkey CD196/CCR6 Antibody[G034E3] (left)

or Elab Fluor® 700 Mouse IgG2b, κ Isotype Control (right).  
Cells in the lymphocytes gate were used for analysis.

### Preparation & Storage

|                 |   |
|-----------------|---|
| <b>Storage</b>  | Keep as concentrated solution.<br>This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze. |
| <b>Shipping</b> | Ice bag   |

### Antigen Information

|                        |                                       |
|------------------------|---------------------------------------|
| <b>Alternate Names</b> | CMKBR6;GPR29;STRL22;CKRL3;GPCRY4;CCR6 |
| <b>Uniprot ID</b>      | P51684                                |
| <b>Gene ID</b>         | 1235                                  |

### For Research Use Only

## Background

CCR6, also known as CD196/CCR6, is a chemokine receptor that is expressed on immature dendritic cells, B lymphocytes, and memory T cells. CCR6 binds CCL20, although members of the  $\beta$  defensin family also bind CCR6 with a lower affinity. CCR6 positive cells, and its ligand CCL20, have been detected in numerous organs, especially the secondary lymphoid organ. CCL20 is selectively made by the follicle-associated epithelium (FAE) overlying Peyer's Patches (PPs) and isolated lymphoid follicles (ILFs). CCL20 contributes to the recruitment of CCR6-expressing B cells to these structures. In humans, CCR6 can function to mediate arrest of T cells on dermal endothelial cells and is highly expressed on T cells resident in both normal and psoriatic skin. CCR6 and/or CCL20 have been implicated in the pathogenesis of rheumatoid arthritis and inflammatory bowel disease. Human T cells that are able to produce IL-17 express CCR6. It suggests that CCL20 and CCR6 have a role in inflammatory diseases by recruiting Th17 cells to target tissues.