A Reliable Research Partner in Life Science and Medicine

# Recombinant Human CROT Protein (474 Leu/Val, His Tag)

Catalog Number: PKSH031306

Note: Centrifuge before opening to ensure complete recovery of vial contents.

# Description

Species Human

Source Baculovirus-Insect Cells-derived Human CROT protein Met 1-Leu 612, 474 Leu/Val,

with an C-terminal His

 Mol\_Mass
 71.5 kDa

 Accession
 Q9UKG9

**Bio-activity** Not validated for activity

#### **Properties**

**Purity** > 93 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per  $\mu$ g of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile 50mM Tris, 100mM NaCl, pH 8.0, 10% glycerol


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

# Data



> 93 % as determined by reducing SDS-PAGE.

# Background

Carnitine octanoyltransferase (CROT or COT), also known as octanoyl-CoA: L-carnitine O-octanoyltransferase, medium-chain/long-chain carnitine acyltransferase, and carnitine medium-chain acyltransferase, is a carnitine acyltransferase belonging to the family of transferases, specifically those acyltransferases transferring groups other than aminoacyl groups that catalyzes the reversible transfer of fatty acyl groups between CoA and carnitine. Carnitine octanoyltransferase (CROT or COT) facilitate the transport of medium- and long-chain fatty acids through the peroxisomal and mitochondrial membranes. It is physiologically inhibited by malonyl-CoA. COT also has functions in efficiently converting one of the end products of the peroxisomal beta-oxidation of pristanic acid, 4, 8-dimethylnonanoyl-CoA, to its corresponding carnitine ester.

### For Research Use Only

Fax: 1-832-243-6017