

A Reliable Research Partner in Life Science and Medicine

Elab Fluor® 647 Anti-Mouse CD103 Antibody[M290]

Catalog Number: E-AB-F1090UM

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Host Rat

Isotype Rat IgG2a, κ **Clone No.** M290

Isotype Control Elab Fluor® 647 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833M]

Conjugation Elab Fluor® 647

Conjugation Information Elab Fluor[®] 647 is designed to be excited by the Red laser (627-640 nm) and detected

using an optical filter centered near 670 nm (e.g., a 660/20 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/10⁶ cells in 100 μ L volume].

Data



C57BL/6 murine splenocytes are stained with Elab Fluor[®] 647 Anti-Mouse CD103 Antibody and Elab Fluor[®] 488 Anti-Mouse CD3 Antibody (Left). Splenocytes stained with Elab

Fluor[®] 488 Anti-Mouse CD3 Antibody (Right) are used as control.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD103;Integrin alpha M290;Integrin alpha-E;Itgae

 Uniprot ID
 Q60677

 Gene ID
 16407

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Background

CD103 is a type I transmembrane glycoprotein known as αE integrin or Integrin αIEL chain. It belongs to the integrin family and is primarily found on intestinal intraepithelial lymphocytes (IEL). CD103 is also expressed on a subpopulation of lamina propria T cells, epithelial dendritic cells, lamina propria-derived dendritic cells, and a small subset of peripheral lymphocytes. T regulatory cells express high level of CD103. The CD103 expression on lymphocytes can be induced upon activation and TGF- β stimulation. In association with integrin $\beta 7$, CD103 is expressed as $\alpha E/\beta 7$ heterodimer. Mature CD103 protein can be cleaved into 2 chains, a 150 kD (C-terminal) chain and a 25 kD (N-terminal) chain, which remain linked by disulfide bonds. CD103 binds to E-cadherin and mediates homing of lymphocytes to the intestinal epithelium.

Fax: 1-832-243-6017