

Recombinant Mouse DPP7/DPPII/DPP2 Protein (His Tag)

Catalog Number: PKSM040386

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse DPP7/DPPII/DPP2 protein Met1-Arg 506, with an C-terminal His
Calculated MW	53.7 kDa
Observed MW	65 kDa
Accession	Q9ET221
Bio-activity	Measured by its ability to cleave the fluorogenic peptide substrate, Lys-Pro-AMC(KP-AMC). The specific activity is > 20, 000 pmoles/min/μg.

Properties

Purity	> 92 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 92 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

DPP7 (dipeptidylpeptidase 7), also known as DPPII and DPP2, is a post-proline cleaving aminopeptidase expressed in quiescent lymphocytes. Dipeptidyl peptidases (DPPs) have post-proline dipeptidyl aminopeptidase activity, cleaving Xaa-Pro dipeptides from the N-termini of proteins. DPPs mediate regulatory activity of their substrates and have been linked to a variety of diseases including type 2 diabetes, obesity and cancer. DPPs can bind specific voltage-gated potassium channels and alter their expression and biophysical properties and may also influence T cells. DPP proteins include DPRP1, DPRP2, DPP3, DPP7, DPP10, DPPX and CD26. It localizes to lysosomes. DPP7 localizes to lysosomes and exists as a homodimer via its leucine zipper motif and is involved in the degradation of oligopeptides. In response to calcium release, it can be secreted in its active form. It is essential for lymphocyte survival, as the inhibition of DPP7 results in quiescent cell apoptosis.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017