AF/LE Purified Anti-Human CD40 Antibody[G28.5]

catalog number: E-AB-F12140



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human CD40 protein

Host Mouse

Isotype Mouse IgGl, κ

Clone G28.5

Purification >98%, Protein A/G purified


Conjugation None (AF/LE)

buffer Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method

.

| Applications | Recommended Dilution |
|--------------|---|
| FCM | $2 \mu g/mL(1\times10^5-5\times10^5 \text{ cells})$ |

Data



Human peripheral blood lymphocytes were stained with 0.2 μg AF/LE Purified Anti-Human CD40 Antibody[G28.5] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD19 PE-conjugated Monoclonal Antibody.

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles. This preparation contains no preservatives, thus it should be handled

under aseptic conditions.

Shipping Ice bag

Background

CD40 is a 48 kD type I glycoprotein also known as BP50. It is a member of the TNFR superfamily primarily expressed on B cells, macrophages, follicular dendritic cells, endothelial cells, fibroblasts, and at low levels on plasma cells. CD40 has been reported to be involved in B cell differentiation, costimulation, isotype class-switching, and protection of B cells from apoptosis. Additionally, CD40 is important for T cell-B cell interactions. The ligand of CD40 is CD154 (CD40 ligand)

For Research Use Only