



A Reliable Research Partner in Life Science and Medicine

# Monkey ANG1 Antibody Pair Set

Catalog No. E-KAB-0666 Applications ELISA

**Synonyms** ANG-1;ANGPT1;AGP1;AGPT

## Kit components & Storage

Title	Specifications	Storage
Monkey ANG1 Capture Antibody	1 vial, 100 µ g	Store at -20°C for one year. Avoid
		freeze/thaw cycles.
Monkey ANG1 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year. Avoid
(Biotin)		freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0666)	
		Monkey ANG1 Capture Antibody	Monkey ANG1 Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Monkey ANG1 protein	Recombinant Monkey ANG1 protein
Information	Swissprot	A0A2K5VW43	
Product details	Reactivity	Monkey	Monkey
	Host	Mouse	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%
		glycerol; pH 7.5	protective protein; 50% glycerol; pH
			7.5
	Purify	Protein A or G	Antigen Affinity
	Specificity	Detects Monkey ANG1 in ELISAs.	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>



A Reliable Research Partner in Life Science and Medicine

### **Applications**

Monkey ANG1 Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Monkey ANG1 Capture	
Capture		Antibody	10
			8 I
			Optical Density
ELISA	1:1000-1:10000	Monkey ANG1 Detection	O O O O O O O O O O O O O O O O O O O
Detection		Antibody (Biotin)	· ·
			0.01
			Monkey ANG1 Concentration (pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

## **Background**

Angiopoietins are proteins with important roles in vascular development and angiogenesis. All angiopoietins bind with similar affinity to an endothelial cell-specific tyrosine-protein kinase receptor. The protein encoded by this gene is a secreted glycoprotein that activates the receptor by inducing its tyrosine phosphorylation. It plays a critical role in mediating reciprocal interactions between the endothelium and surrounding matrix and mesenchyme and inhibits endothelial permeability. The protein also contributes to blood vessel maturation and stability, and may be involved in early development of the heart. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>

Tel: 1-832-243-6086 Fax: 1-832-243-6017