

Recombinant CD3 Monoclonal Antibody

catalog number: **AN301033L**

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human;Mouse;Rat
Immunogen	Recombinant Human CD3 protein
Host	Rabbit
Isotype	IgG, κ
Clone	5A8
Purification	Protein A
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications


Applications	Recommended Dilution
IHC	1:200-1:1000
WB	1:1000-1:5000

Data




Western Blot with Recombinant CD3 Monoclonal Antibody at dilution of 1:1000. Lane A: Jurkat cells.

Observed-MW:23 kDa
Calculated-MW:23 kDa



Immunohistochemistry of paraffin-embedded mouse spleen tissue using Recombinant CD3 Monoclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded human tonsil tissue using Recombinant CD3 Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.0

The protein encoded by this gene is the CD3-epsilon polypeptide, which together with CD3-gamma, -delta and -zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T-cell receptor-CD3 complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cell development. Defects in this gene cause immunodeficiency. This gene has also been linked to a susceptibility to type I diabetes in women.