

## Elab Fluor® 647 Anti-Human CD55 Antibody[HI55a]

Catalog Number: AN00336M

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Isotype</b>	Mouse IgG2a, κ
<b>Clone No.</b>	HI55a
<b>Isotype Control</b>	Elab Fluor® 647 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802M]
<b>Conjugation</b>	Elab Fluor® 647
<b>Conjugation Information</b>	Elab Fluor® 647 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 670 nm (e.g., a 660/20 nm bandpass filter).
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

### Applications

### Recommended usage

<b>FCM</b>	Each lot of this antibody is quality control tested by flow cytometric analysis. <b>The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).</b> Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	---

### Data



Staining of normal human peripheral blood cells with Elab Fluor® 647 Anti-Human CD55 Antibody[HI55a] (filled gray histogram) or Elab Fluor® 647 Mouse IgG2a, κ Isotype Control (empty black histogram). Cells in the lymphocytes gate were used for analysis.

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	Lyb-2;Ly-19.2;Ly-32.2
<b>Uniprot ID</b>	P13591
<b>Gene ID</b>	1604

### For Research Use Only

## Background

CD55 is a 60-70 kD glycosylphosphatidylinositol (GPI)-anchored single chain glycoprotein also known as decay-accelerating factor (DAF). It is expressed on hematopoietic cells including erythrocytes and many non-hematopoietic cells. CD55 accelerates the dissociation of the components of the C3-convertases (namely C2a from C4b in the C4bC2a complex, a C3-convertase of the classical pathway, and factor Bb from the C3bBb complex, a C3-convertase of the alternative pathway) to protect cells from inappropriate damage caused by autologous complement. CD55 has been reported to reduce the efficiency of NK cell lysis and induce signal transduction in T cells. CD55 has also been shown to interact with CD97 and bind to Coxsackie and Echovirus.