

A Reliable Research Partner in Life Science and Medicine

Recombinant CDKN2A/p14ARF Monoclonal Antibody

catalog number: AN301615L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human;

Immunogen Recombinant human CDKN2A/p14ARF fragment


HostRabbitIsotypeIgG, κCloneA318


Purification Protein A purified

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution WB 1:500-1:2000 IHC 1:200-1:1000 IF 1:50 IP 1:50-1:100


Data






Western Blot with CDKN2A/p14ARF Monoclonal Antibody at Immunohistochemistry of paraffin-embedded Human lung dilution of 1:2000. Lane 1: HeLa squamous carcinoma using CDKN2A/p14ARF Monoclonal Observed-MW:14 kDa Antibody at dilution of 1:1000.

Observed-MW:14 kDa Calculated-MW:14 kDa





Immunofluorescent analysis of (4% Paraformaldehyde) fixed Immunoprecipitation analysis using anti-CDKN2A/p14ARF

HeLa cells using anti-CDKN2A/p14ARF Monoclonal Monoclonal Antibody. Western blot was performed from the immunoprecipitate using CDKN2A/p14ARF Monoclonal

Immunoprecipitation analysis using anti-CDKN2A/p14ARF
Monoclonal Antibody. Western blot was performed from the
immunoprecipitate using CDKN2A/p14ARF Monoclonal
Antibody at a dilution of 1:100. Lane 1: 5% Input, Lane 2:
CDKN2A/p14ARF Monoclonal Antibody, Lane 3: Rabbit
monoclonal IgG Isotype

Observed-MW:14 kDa
Calculated-MW:14 kDa

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com



Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Shipping Ice bag

Background

Capable of inducing cell cycle arrest in G1 and G2 phases. Acts as a tumor suppressor. Binds to MDM2 and blocks its nucleocytoplasmic shuttling by sequestering it in the nucleolus. This inhibits the oncogenic action of MDM2 by blocking MDM2-induced degradation of p53 and enhancing p53-dependent transactivation and apoptosis. Also induces G2 arrest and apoptosis in a p53-independent manner by preventing the activation of cyclin B1/CDC2 complexes. Binds to BCL6 and down-regulates BCL6-induced transcriptional repression.

For Research Use Only