# **Elabscience**®

### **Porcine IL-6 Antibody Pair Set**

Catalog No.	E-KAB-0612	Applications	ELISA
Synonyms	IL16;LCF;NIL16;PRIL16;PrIL-16		

#### **Kit components & Storage**

Title	Specifications	Storage
Porcine IL-6 Capture Antibody	1 vial, 100 µ g	Store at -20°C for one year. Avoid
		freeze/thaw cycles.
Porcine IL-6 Detection Antibody (Biotin)	1 vial, 50 μL	Store at -20°C for one year. Avoid
		freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0612)		
		Porcine IL-6 Capture Antibody	Porcine IL-6 Detection Antibody (Biotin)	
Immunogen	Immunogen	Recombinant Porcine IL-6 protien	Recombinant Porcine IL-6 protien	
Information	Swissprot	P26893		
Product details	Reactivity	Porcine	Porcine	
	Host	Goat	Goat	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5 mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%	
		glycerol; pH 7.5	protective protein; 50% glycerol; pH	
			7.5	
	Purify	Antigen Affinity	Antigen Affinity	
	Specificity	Detects Porcine IL-6 in ELISAs.		

For Research Use Only

## **Elabscience**®

#### Applications

Porcine IL-6 Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Porcine IL-6 Capture	
Capture		Antibody	10
			Optical Density
ELISA	1:1000-1:10000	Porcine IL-6 Detection	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
Detection		Antibody (Biotin)	
			0.01 1 10 100 1000
			Porcine IL-6 Concentration (pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

#### Background

Cytokine with a wide variety of biological functions. It is a potent inducer of the acute phase response. Plays an essential role in the final differentiation of B-cells into Ig-secreting cells Involved in lymphocyte and monocyte differentiation. It induces myeloma and plasmacytoma growth and induces nerve cells differentiation Acts on B-cells, hepatocytes, hematopoeitic progenitor cells and cells of the CNS. Also acts as a myokine. It is discharged into the bloodstream after muscle contraction and acts to increase the breakdown of fats and to improve insulin resistance.

For Research Use Only