A Reliable Research Partner in Life Science and Medicine

AMACR Monoclonal Antibody

catalog number: E-AB-22010

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat **Immunogen** Synthetic Peptide

HostMouseIsotypeIgGClone1F1


Purification Protein A purification

Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein

protectant and 50% glycerol.


Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:100-1:300
IF	1:100-1:300

Data



Western Blot analysis of 1) HepG2, 2) Mouse kidney using AMACR Monoclonal Antibody at dilution of 1:1000.

Observed-MV:42 kDa Calculated-MV:42 kDa



Negative Control

mmunohistochemistry of paraffin-embedded

Immunohistochemistry of paraffin-embedded Human stomach tissue using AMACR Monoclonal Antibody at dilution of 1:200.

Immunofluorescence analysis of Mouse kidney tissue using AMACR Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Tel: 1-832-243-6086 Email:techsupport@elabscience.com

Elal

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Elabscience®

This gene encodes a racemase. The encoded enzyme interconverts pristanoyl-CoA and C27-bile acylCoAs between their (R)- and (S)-stereoisomers. The conversion to the (S)-stereoisomers is necessary for degradation of these substrates by peroxisomal beta-oxidation. Encoded proteins from this locus localize to both mitochondria and peroxisomes. Mutations in this gene may be associated with adult-onset sensorimotor neuropathy, pigmentary retinopathy, and adrenomyeloneuropathy due to defects in bile acid synthesis. Alternatively spliced transcript variants have been described. Read-through transcription also exists between this gene and the upstream neighboring C1QTNF3 (C1q and tumor necrosis factor related protein 3) gene.

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Tel: 1-832-243-6086 Email:techsupport@elabscience.com

Rev. V1.6