Recombinant Axin 2 Monoclonal Antibody


catalog number: E-AB-81534

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|---|
| Reactivity | Human;Mouse;Rat |
| Immunogen | A synthetic peptide of human Axin 2 |
| Host | Rabbit |
| Isotype | IgG |
| Clone | R06-3A8 |
| Purification | Affinity Purified |
| Buffer | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.05% stabilizer and 0.05% |
| | protective protein. |
| | |

| Applications | Recommended Dilution |
|--------------|-----------------------------|
| WB | 1:500-1:1000 |

Data



Western blot detection of Axin 2 in K562,C6,3T3,Hela cell lysates using Axin 2 Rabbit mAb(1:1000 diluted).Predicted band size:94kDa.Observed band size:94kDa.

> Observed-MW:94 kDa Calculated-MW:94 kDa

| Preparation & Storage | |
|-----------------------|---|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the |
| | temperature recommended. |

Background

The Axin-related protein, Axin2, presumably plays an important role in the regulation of the stability of beta-catenin in the Wnt signaling pathway, like its rodent homologs, mouse conductin/rat axil. In mouse, conductin organizes a multiprotein complex of APC (adenomatous polyposis of the colon), beta-catenin, glycogen synthase kinase 3-beta, and conductin, which leads to the degradation of beta-catenin. Apparently, the deregulation of beta-catenin is an important event in the genesis of a number of malignancies. The AXIN2 gene has been mapped to 17q23-q24, a region that shows frequent loss of heterozygosity in breast cancer, neuroblastoma, and other tumors. Mutations in this gene have been associated with colorectal cancer with defective mismatch repair.

For Research Use Only