Recombinant Human OSMR/IL31RB Protein (aa 28-739, His Tag)

Catalog Number: PKSH032836



Note: Centrifuge before opening to ensure complete recovery of vial contents. Description **Species** Human Mol Mass 82.0 kDa Accession O99650 **Bio-activity** Not validated for activity **Properties** > 90 % as determined by reducing SDS-PAGE. Purity < 1.0 EU per µg of the protein as determined by the LAL method. Endotoxin Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 Storage °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. This product is provided as lyophilized powder which is shipped with ice packs. Shipping Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2. Formulation Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. Reconstitution Please refer to the printed manual for detailed information.

	•
kDa	MK
120	Elabs. An and
90	
60	Tabscienc
40	E
30	
-tabsciel	
40 30 20	
14	

> 90 % as determined by reducing SDS-PAGE.

Background

Data

Oncostatin-M-Specific Receptor Subunit β (OSMR β) is a 150 - 180 kDa member of the IL-6 receptor family. OSMR β associates with gp130 to form the type II OSM receptor, the receptor is responsive to OSM. Gp130 subunit is shared by other IL-6 family cytokine receptors, and OSMR β associates with gp130-like receptor (GPL) to form a receptor complex responsive to IL-31. The human OSMR β cDNA encodes a 979 amino acid (aa) precursor, the precursor includes a 27 aa signal sequence, a 712 aa extracellular domain (ECD), a 22 aatransmembrane segment, and a 218 aa cytoplasmic domain. The ECD contains one partial and one complete hematopoietin domain, an Ig-like domain, and three Fibronectin type-III domains.

For Research Use Only