## **Elabscience**®

## Purified Anti-Human CD3 Antibody[OKT-3]

catalog number: E-AB-F1001A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

-	e opening to ensure complete recovery of var contents.
Description	
Reactivity	Human
Immunogen	Recombinant Human CD3 protein
Host	Mouse
Isotype	Mouse IgG2a, κ
Clone	OKT3
Purification	>98%, Protein A/G purified
Conjugation	Unconjugated
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze
	to completely remove the stabilizer prior to labeling.
Applications	Recommended Dilution
FCM	$2 \ \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$
Data	
Court 0 20 Court 0 101	10 <sup>2</sup> 10 <sup>3</sup> 10 <sup>4</sup> 10 <sup>5</sup> 10 <sup>6</sup> 10 <sup>7</sup>
Human peripheral bloc	od lymphocytes were stained with 0.2
μg Purified Anti-Huma	an CD3 Antibody[OKT-3](Right) and
0.2 µg Mouse IgG2a,	κ Isotype Control(Left), followed by
PE-conjugated Goat A	Anti-Mouse IgG Secondary Antibody.
Preparation & Storag	e
Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

i reparation & Storage	
Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /
	thaw cycles.
Shipping	Ice bag
Background	

CD3 $\epsilon$  is a 20 kD chain of the CD3/T cell receptor (TCR) complex, which is composed of two CD3 $\epsilon$ , one CD3 $\gamma$ , one CD3 $\delta$ , one CD3 $\zeta$  (CD247), and a T cell receptor ( $\alpha/\beta$  or  $\gamma/\delta$ ) heterodimer. It is found on all mature T lymphocytes, NK T cells, and some thymocytes. CD3, also known as T3, is a member of the immunoglobulin superfamily that plays a role in antigen recognition, signal transduction, and T cell activation.